

BOER & PYGMY GOAT SHOTS

MOMS

2-4 WEEKS BEFORE BIRTH

2cc Covexin 8 (sq)

BoSe - 1 cc per 40# (mus)

1cc J-5 E-Coli (sq)

BABIES AFTER BIRTH

Iodine navels with weak Iodine

1/4 cc BoSe (mus)

1/2 tube Coliumune Oral

1/4 cc A&D (mus)

BABIES 2 WEEKS OLD RIGHT BEFORE CAST/DEHORN

1/2 bottle of Tetanus (sq)

2 cc C & D Antitoxin (sq)

1 cc CL (mus)

Weigh

BABIES 6 WEEKS OLD

1 cc CL (mus)

Weigh

BABIES 2 MONTHS OLD

2cc CD/T (sq)

2cc BSRV (intranasal)

2cc Pasteurella (sq)

Worm

BABIES 3 MONTHS OLD

2cc A&D (mus)

2cc CD/T (sq)

2cc BSRV (intranasal)

2cc Pasteurella (sq) (booster)

Worm

Weigh

BABIES 4 MONTHS OLD

2cc Triangle (sq)

Worm

BABIES 5 MONTHS OLD

2cc Triangle (sq) (booster)

BABIES 6 MONTHS OLD

1cc per 40# BoSe (mus)

2cc CD/T (sq)

Worm

Weigh

BABIES 8 & 10 MONTHS OLD

Worm

BABIES 1 YEAR OLD

1cc per 40# BoSe (mus)

2cc Covexin 8 (sq)

2cc A&D (mus)

2cc Pasteurella (sq)

Worm

Weigh

MOM THAT DID NOT GET E-COLI LAST YEAR:

1cc 6 weeks before birth and then 1cc 3 weeks

before birth - next season just 1cc 3 weeks

before birth.

NON BREEDING DOES, BUCKS, WETHERS

FALL SHOTS

2cc BoSe (mus)

2cc Covexin 8 (sq)

2cc A&D (mus)

2cc Triangle (sq)

ALL GOATS OVER 1 YEAR

2cc Pasteurella JAN & JULY

WORM JAN, APRIL, JULY, OCT THIS IS FOR ANY GOAT OVER 1 YEAR

SUGGESTED WORMERS:

Ivomec Drench (orally) - external bugs OK FOR PREGNANT DOES

Cydetin (orally) - external bugs OK FOR PREGNANT DOES

Dectomax - Injection

Valbazen (orally) - Tape Worms, not external bugs

DO NOT USE ON PREGNANT DOES

Safeguard/Panacure - Drench OK FOR PREGNANT DOES

(sq) = under skin

(mus) = in the muscle

The above listed shots is followed by Newton Farms, Lakeville Indiana for all Boer & Pygmy goats.

REVISED 1/27/11

www.newtonfarms.org